



BMA BIOMEDICALS

BMA BIOMEDICALS
Rheinstrasse 28-32
CH-4302 Augst (Switzerland)
Phone: ++41 61 811 6222
Fax: ++41 61 811 6006
info@bma.ch
www.bma.ch

Monoclonal Antibody To Mouse CD117 (c-kit) Receptor for Stem Cell Factor

Monoclonal antibody ACK4 recognizes the receptor for Stem Cell Factor (SCF) / Steel Factor. The cellular function of CD117 is that of a growth factor receptor. The ligand exists in both soluble and membrane form. The interaction between SCF and c-kit is essential for the development of hematopoietic, gonadal and pigment stem cells. c-kit positive cells are a subset of CD34⁺ hematopoietic precursor cells, expression of CD117 can be found on 5-10% of total adult bone marrow cells.

Product Number:	T-2033 (Lot 01PO1004)
Clone:	ACK4
Host species, isotype:	Rat IgG2a
Quantity:	200µg
Format:	Affinity purified, lyophilized Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.4mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 5mg/ml bovine serum albumin (BSA) as a stabilizer and 0.09% sodium azide as a preservative.
Stability:	Original vial: 1 year at 4° - 8°C Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.
Applications:	Tested for immunohistochemistry (IHC); has been described to work in FACS. Approximate working dilution for IHC: Frozen sections: 0.4µg/ml (1:500) Paraffin sections: does not react on routinely processed paraffin sections. Suggested positive control: Mouse spleen and tongue Please see www.bma.ch for protocols and general information.
Immunogen:	IL-3 dependent mast cells derived from WB+/+ mice.
Antigen, epitope:	The antigen is a 145kD membrane protein (reduced or unreduced) that belongs to the immunoglobulin supergene family. The epitope has not been characterized.

Antigen distribution: **Isolated Cells:** The antigen is described to be present on 5-10% of total adult bone marrow cells.
Tissue Sections: The antigen is expressed on CD117 positive cells in various tissues e.g. in mouse spleen and tongue. There is a staining on cells in the red pulp of spleen, whereas there are only few cells with strong staining in tongue.

Specificity: **Mouse:** receptor for stem cell factor
Other species: not tested

Selected references

Ogawa, M. et al.: Expression and Function of c-kit in Hematopoietic Progenitor Cells. J. Exp. Med 174: 63-71 (1991)
Suda, T. et al.: A Stimulatory effect of Recombinant Murine Interleukin-7 (IL-7) on B-Cell Colony Formation and Inhibitory Effect of IL-1 α . Blood 74(6): 1936-41 (1989).
Kunisada, T. et al.: Murine Cutaneous Mastocytosis and Epidermal Melanocytosis Induced by Keratinocyte Expression of Transgenic Stem Cell Factor: J. Exp. Med. 187(10): 1565-73 (1998).

For *in vitro* research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.