BMA BIOMEDICALS

Monoclonal Antibody To Human CD120b

Anti human Tumor Necrosis Factor (TNF) - Receptor p75

Monoclonal antibody utr 1 is useful for studying biological effects of TNF-receptor p75 *in vitro* where it inhibits the binding of radiolabeled TNF to human cells expressing the p75 TNF receptor. In order to completely block TNF binding to the cell surface, 10µg/ml of utr 1 are required. utr 1 itself may have an agonistic effect in assays measuring cytotoxicity, fibroblast growth or IL-6 secretion. See also Product T-1409, anti human CD120a (TNF-receptor p55), clone htr 9.

This antibody was produced serum-free, without fetal calf serum. Packaged under nitrogen.

TECHNICAL AND ANALYTICAL CHARACTERISTICS

Product Number:	T-1412 (Lot 10PO1607)
Clone:	utr 1
Host species, isotype:	Mouse IgG1
Quantity:	500µg
Format:	Affinity purified, lyophilizer Reconstitute by adding 0.5ml distilled water. This stock solution contains 1mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), no stabilizer and no preservative.
Stability:	Original vial: 6 months at 4° - 8°C from date of purchase.
Applications:	Tested for immunohistochemistry (IHC); has also been described to work in FACS, Western Blots, immunoprecipitation. **Approximate working dilution for IHC:** Frozen sections: 4µg/ml (1:250) Paraffin section:20µg/ml (1:50); pretreatment with Proteinase K for antigen retrieval is recommended Optimal dilutions should be determined by the end user Please see **www.bma.ch** for protocols and general information.
Immunogen:	Partially purified TNF binding proteins
Protein database:	human-P20333

©2024 BMA BIOMEDICALS, Rheinstrasse 28-32, CH-4302 Augst (Switzerland)
Phone: +41 61 811 6222, Email: info@bma.ch, www.bma.ch

Antigen distribution:	**Isolated cells:** Isolated cells and cell lines: U937, HL-60. Lymphocytes from peripheral blood show a faint staining. Mitogen stimulation of lymphocytes increases the intensity. Bone marrow cells are not stained with utr 1.
Tissue sections:	Immunohistochemical staining in normal healthy tissue is confined to the lymphohistiocytic tissue, which includes the thymus and lymphoid organs such as spleen, tonsils, lymph nodes, mucosa, and associated lymphoid tissue (see ref 7). Expression of CD120a and CD120b can be detected in different areas where an overlapping is found between CD120b and IL-2 receptor expression. CD120b expression can be detected mainly in the T-cell area whereas CD120a expression is restricted to dendritic reticulum cells in the germinal centres. In non-lymphoid organs (kidney, liver, heart, brain, adrenals, uterus, ovary, testes, prostate, stomach, intestines) CD120b recognition is restricted to some interstitial reticulum cells in the kidney. Cells which are known to respond to TNF namely endothelial cells, smooth muscle cells and fibroblasts did not show expression of CD120a or CD120b. Investigations on pathological tissues show a CD120b expression on epitheloid cell granuloma and giant cells in sarcoidosis.
Specificity:	**Human:** CD120b **Other species:** not tested

Selected references

Brockhaus, M., et al.: Identification of two types of necrosis factor receptors on human cell lines by monoclonal antibodies. Proc. Natl. Acad Sci USA (1990) **87**: 3127-3131

Espevik, T.,et al.: Characterization of binding and biological effects of monoclonal antibodies against a human tumor necrosis factor receptor. J Exp Med (1990) **171**:415-426

Hohmann, H., et al.: Expression of the type A and B Tumor Necrosis Factor (TNF) receptors is independently regulated and both receptors mediate activation of the transcription factor NF-κB. J Biol Chem (1990) **265**: 22409-22417

Shalaby, M.R., et al.: Binding and regulation of cellular functions by monoclonal antibodies against human tumor necrosis factor receptors. J Exp Med (1990) **172**: 1517-1520

Gehr, G., et al.: Both tumor necrosis factor receptor types mediate proliferative signals in human mononuclear cell activation. J Immunol (1992) **149**: 911-917

Vandenabeele, P., et al.: Functional characterization of the human tumor necrosis factor receptor p75 in a transfected rat/mouse T cell hybridoma. J Exp Med (1992) **176**: 1015-1024

Ryffel B., et al.: TNF receptors in lymphoid tissues and lymphomas: Source and site of action of tumor necrosis factor alpha. Am J Pathol (1991) 139: 7-15.

For in vitro research only. This product contains no preservative.

©2024 BMA BIOMEDICALS, Rheinstrasse 28-32, CH-4302 Augst (Switzerland)
Phone: +41 61 811 6222, Email: info@bma.ch, www.bma.ch