



BMA BIOMEDICALS



Peninsula Laboratories

Rabbit anti Endothelin-1

Endothelin (ET-1) is a 21 amino acid peptide, also known as preproendothelin-1 (PPET1), that is produced by the vascular endothelium from a 39 amino acid precursor, big ET-1, through the actions of an endothelin converting enzyme (ECE) found on the endothelial cell membrane. ET-1 is involved in regulation of cardiovascular functions where it has been implicated in the pathogenesis of hypertension, coronary vasospasm, and heart failure. It is also known to cause fibrosis of the vascular cells as well as to stimulate production of reactive oxygen species.

This antibody was generated by immunization of rabbits with Endothelin-1 coupled to a carrier protein.

TECHNICAL AND ANALYTICAL CHARACTERISTICS

Catalog Number:	T-4050 (Lot A18PO23091)
Clonality:	Polyclonal
Host species:	Rabbit
Reactivity:	Human, Bovine, Dog, Mouse, Porcine, Rat
Quantity:	50 µl
Format:	Neat undiluted antiserum, lyophilized, packaged under nitrogen. Reconstitute by adding 50 µl distilled water. This will give the equivalent of undiluted antiserum.
Stability and Storage:	Original vial: 1 year at 4° - 8°C from date of purchase. Stock solution and aliquots thereof: 1 year at -20°C. Avoid freeze-thaw cycles.
Immunogen:	Synthetic peptide H-Cys-Ser-Cys-Ser-Ser-Leu-Met-Asp-Lys-Glu-Cys-Val-Tyr-Phe-Cys-His-Leu-Asp-Ile-Ile-Trp-OH, (Disulfide bonds between Cys1 and Cys15/Cys3 and Cys11 coupled to carrier protein.
Protein database:	n/a

CROSS REACTION:

PEPTIDE	%
Endothelin-1 (human, bovine, dog, mouse, porcine, rat)	100
Endothelin-2 (human, canine)	100
Big Endothelin-1 (human)	100
Big Endothelin-1 (porcine)	100
Endothelin-3 (human, rat, porcine, rabbit)	0
BNP-32 (human)	0

CROSS REACTION:

PEPTIDE	%
ANF (1-28) (human)	0
Angiotensin I	0
Angiotensin II	0
[Arg ⁸]-Vasopressin	0
VIP (human, bovine, porcine, rat)	0
Ac- β -Endorphin (human)	70
Ac- β -Endorphin (1-27) (human)	50
β -Endorphin (6-31) (human)	43

Applications:

Tested in immunohistochemistry (IHC) and ELISA against Endothelin-1. Other applications like FACS or Western Blot may work as well.

Optimal dilutions should be determined by the end user.

Please see **bma.ch** for protocols and general information.

Approximate working dilution for IHC:

Frozen sections: not tested yet.

Paraffin sections: 1:500; recommended Proteinase K pre-treatment.

Related Products:

T-4048: anti Endothelin-1, diluted antiserum, host: rabbit

T-4049: anti Endothelin-1, purified IgG

T-5011: anti Endothelin-1, neat antiserum, host: guinea pig

T-5010: anti Endothelin-1, diluted antiserum, host: guinea pig

S-1156: ELISA, extraction-free for serum and plasma samples

This product contains no preservative and is intended for laboratory use and research purposes only. Not for diagnostic or therapeutic applications.