



BMA BIOMEDICALS



Peninsula Laboratories

Rabbit anti Amylin

Amylin (also known as islet amyloid polypeptide (IAPP)) is a 37 amino acid peptide hormone with a disulfide bridge between Cys² and Cys⁷. Both the amidated C-terminus and the disulfide bridge are necessary for the full biological activity of Amylin. It is co-secreted with insulin from the pancreatic β -cells and plays an important role in the glycemic control, regulating it by slowing gastric emptying and promoting satiety. It is also involved in bone metabolism, along with the related peptides calcitonin and calcitonin gene related peptide.

This antibody was generated by immunization of rabbits with Amylin coupled to a carrier protein.

TECHNICAL AND ANALYTICAL CHARACTERISTICS

Catalog Number:	T-4149 (Lot A18PO2147)
Clonality:	Polyclonal
Host species:	Rabbit IgG
Reactivity:	Human
Quantity:	400 μ g
Format:	Protein A affinity purified from antiserum, lyophilized. This stock solution contains 2mg/ml IgG, phosphate buffered saline pH 7.4 (PBS), 6mg/ml BSA as a stabilizer, and 0.05% (w/v) ProClin TM 150 as a preservative.

Reconstitute by adding 0.2ml of distilled water.

Stability and Storage: Original vial: 1 year at 4° - 8°C from date of purchase.
Stock solution and aliquots thereof: 1 year at -20°C. Avoid freeze-thaw cycles.

Immunogen: Synthetic peptide H-Lys-Cys-Asn-Thr-Ala-Thr-Cys-Ala-Thr-Gln-Arg-Leu-Ala-Asn-Phe-Leu-Val-His-Ser-Ser-Asn-Asn-Phe-Gly-Ala-Ile-Leu-Ser-Ser-Thr-Asn-Val-Gly-Ser-Asn-Thr-Tyr-NH₂, (disulfide bond) coupled to carrier protein.

Protein database: [human-P10997](#)

Cross-Reactivity:

PEPTIDE	%
Amylin (free acid) (human)	100
Amylin (human)	100
Amylin (rat)	100

Cross-Reactivity:

PEPTIDE	%
Amylin (feline)	100
α -CGRP (rat)	75
α -CGRP (human)	1
Calcitonin (human)	0
NPY (porcine)	0
Pancreastatin (porcine)	0
VIP (human, bovine, rat, porcine)	0
Insulin (human)	0
Glucagon (1-29) (human, bovine, porcine)	0
Somatostatin-14	0
Pancreatic Polypeptide (human)	0
Proinsulin (human)	0

Applications:

Tested in ELISA against Amylin and in immunohistochemistry (IHC). Other applications like FACS and Western blot may work as well.

Approximate working dilution for IHC:

Paraffin sections: (1:800); pretreatment with Proteinase K for antigen retrieval is recommended

Optimal dilutions should be determined by the end user

Please see www.bma.ch for protocols and general information

Related Products:

T-4150: Rabbit anti Amylin, neat antiserum (hu)
T-4148: Rabbit anti Amylin, diluted antiserum (hu)
S-1187: ELISA high sensitivity for extracted samples
S-1420: ELISA extraction-free for human samples
S-1413: ELISA extraction-free for rat samples

This product contains ProClin™ 150 as a preservative and is intended for laboratory use and research purposes only. Not for diagnostic or therapeutic applications.