



Normal Rabbit Serum, pretitrated

This serum has been designed to be used in conjunction with pretitered Goat anti-Rabbit IgG (catalog #T-4576) for precipitation of "bound" counts in the titration of antisera or in RIAs.

TECHNICAL AND ANALYTICAL CHARACTERISTICS

Product number: T-4577 (Lot A19PO24086)

Format: Serum, lyophilized, packaged under nitrogen

Reconstitution Information:

Rehydrate the lyophilized powder with 50 ml of assay buffer.

Recommended Use:

- Add 100µl of rehydrated Goat anti-Rabbit IgG (catalog #T-4576) to each tube
- Add to each tube, 100µl of pretitrated Normal Rabbit Serum, and vortex
- Allow the precipitate to form for 2 hours at room temperature (or 6 hours at 4°C)
- Centrifuge the tube at 1700xg for 20 min to collect the precipitate
- · Collect the precipitate and carefully aspirate the supernatant
- · Count the pellet in a scintillation well gamma counter

Related Products:

T-4576: Goat anti-Rabbit IgG

Y-1000: SEP-Columns (200mg, 50 per box)

Y-1040: Buffer A for sample extraction

Y-1045: Buffer B for sample extraction

Y-1050: RIA Buffer

This product contains no preservative and is intended for laboratory use and research purposes only. Not for diagnostic or therapeutic applications.

T-4577 07.08.2024