
Monoclonal Antibody to human Lambda Light Chain

Marker for B-cells and for Immunoglobulin Isotyping

Monoclonal antibody λ -1 can be used for the detection of surface and intracytoplasmic immunoglobulins on normal and neoplastic B-cells. In conjunction with clone κ -117 (product T-1307), the light chain isotype of a B-cell population can be determined.

Product number: T-1304

Clone: λ -1

Lot: 02PO1313

Host species, isotype: Mouse IgG1, kappa light chain

Quantity: 100 μ g

Format: Affinity purified, lyophilized
Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.2mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 5mg/ml bovine serum albumin (BSA) as a stabilizer and 0.05% sodium azide as a preservative.

Stability: Original vial: 1 year at 4° - 8°C
Stock solution: 1 year at -20°C. Avoid repeated thawing and freezing.

Applications: Tested for immunohistochemistry (IHC)
Approximate working dilution for IHC:
Frozen sections: 0.125 μ g/ml (1:1600)
Paraffin sections: Has been described to work with proteinase K pretreatment.
Optimal dilutions should be determined by the end user.
Suggested positive control: Human tonsil.
Please see www.bma.ch for protocols and general information.

Immunogen: Purified human IgG.

Antigen, epitope: The epitope is located on the lambda light chain of human Ig.

Antigen distribution:

Isolated cells: Antigen receptor bearing B-cells are stained positively.

Tissue sections: λ -1 labels follicular and non-follicular B-cells in lymphoid tissues. Intravascular staining can also be observed caused by circulating Ig. Strong labeling occurs with plasma cells in paraffin embedded tissues.

The antibody stains surface and intracellular immunoglobulins in cryostat sections. Only intracellular light chains are stained in formaldehyde fixed and paraffin embedded tissues after enzyme digestion (eg. plasma cells are stained in the latter case.)

Specificity:

Human: lambda light chain

Other species: not tested.

Selected references

ODERMATT, B.F., KNECHT, H., HAGEN, M.F. FEHR, J., RUETTNER, J.R.: Diagnostic and Prognostic Value of Monoclonal Antibodies in Immunophenotyping of T-cell Lymphomas. Acta Haematologica, 77, 72-77 (1987).

ODERMATT, B.F., MAURER, R., RUETTNER, J.R.: Morphologische und immunologische Kriterien zur Diagnostik der Non-Hodgkin-Lymphome von niedrigem Malignitätsgrad. Schweiz. Rundschau Med. (PRAXIS) 75(15), 401-407 (1986).

STEIN, H., DALLENBACH, F., DIENEMANN, D.: Differentiation of Physiologic and Malignant Cells of the Lymphatic System. Verh. Dtsch. Ges. Path. 72, 57-85 (1988).

For *in vitro* research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.