

---

## **Monoclonal Antibody To Human HLA-DQ**

### **Marker for MHC class II**

Monoclonal antibody FN81 recognizes the HLA-DQ antigen, an HLA class II antigen with homology to murine H-2A. HLA class II antigens are mainly expressed on specialized antigen presenting cells (APCs), dendritic cells, B-cells and macrophages. HLA class II antigen is presented on activated T-cells. HLA class II molecules present exogenously derived antigen to the T-cell receptor (TCR) on CD4<sup>+</sup> T lymphocytes. Both chains in HLA-DQ are polymorphic. HLA-DQ is expressed after DR and DP during haematopoietic progenitor cell development. It is strongly expressed in the thymic cortex.

---

**Product number: T-1360**

**Clone: FN81**

**Lot: 01PO0801**

### **TECHNICAL AND ANALYTICAL CHARACTERISTICS:**

**Host species, subclass:** Mouse IgG2a

**Quantity:** 100µg

**Format:** Affinity purified, lyophilized

Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.2mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 5mg/ml bovine serum albumin (BSA), and 0.09% sodium azide as a preservative.

**Stability:** Original vial: 1 year at 4° - 8°C

Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.

**Applications:** Tested for immunohistochemistry (IHC); has been described to work in FACS.

#### **Approximate working dilution for IHC:**

Frozen sections: ≤0.2µg/ml (1:1000)

Paraffin sections: not tested

Optimal dilutions should be determined by the end user.

Suggested positive control: Human tonsil.

Please see **www.bma.ch** for protocols and general information.

**Immunogen:** Stimulated human B lymphocytes

**Antigen, epitope:** The antigen is HLA-DQ. The epitope has not been further characterized.

<b>Antigen distribution:</b>	<b>Isolated cells:</b> The HLA DQ antigen is present on approximately 10% of peripheral blood lymphocytes.
<b>Specificity:</b>	<b>Human:</b> HLA-DQ <b>Other:</b> not tested.

---

### **Selected references**

Barclay, Brown et al., The Leukocyte Antigen FactsBook, 2nd edition, Harcourt Brace & Company, London, (1997)

Guy, K. et al., (1982). Differential expression and serologically distinct subpopulations of human Ia antigens detected with monoclonal antibodies to Ia alpha and beta chains. Eur. J. Immunol. 12, 942-948.

Symington, F. W. et al., (1985). Differential Ia antigen expression by autologous human erythroid and B lymphoblastoid cell lines. J. Immunol. 135, 1026-1032.

Ishikura, H., Ishikawa, N., Aizawa, M. (1987). Differential expression of HLA-class II antigens in the human thymus. Transplantation 44, 314-317

For *in vitro* research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.