

BMA BIOMEDICALS Rheinstrasse 28-32 CH-4302 Augst (Switzerland) Phone:+41 61 811 6222 Fax: +41 61 811 6006 info@bma.ch www.bma.ch

Monoclonal Antibody to Rat Granulocytes

Monoclonal antibody RK-4 detects all rat granulocytes in frozen sections.

Product number: T-3105 Clone: RK-4 Lot: 03PO1011 TECHNICAL AND ANALYTICAL CHARACTERISTICS: Host species, subclass: Mouse IgG1

nost species, subcidss.	
Quantity:	250µg
Format:	Affinity purified, lyophilized
	Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.5mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 5mg/ml bovine serum albumin (BSA) as a stabilizer and 0.09% sodium azide as a preservative.
Stability:	Original vial: 1 year at 4° - 8°C
	Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.
Applications:	Tested for immunohistochemistry (IHC); has been described to work in FACS.
	Approximate working dilution for IHC: Frozen sections: 1.25µg/ml (1:400) Paraffin sections: does not react on routinely processed paraffin sections. Optimal dilutions should be determined by the end user.
	Suggested positive control: Rat spleen, blood smears.
	Please see www.bma.ch for protocols and general information.
Immunogen:	Peritoneal cells.
Antigen, epitope:	The antigen is localized on the cell surface. The epitope has not been further characterized.
Antigen distribution:	Isolated cells Only a low percentage of cells from peripheral blood stains positive. Positive on peritoneal cells 18h after thioglycollate injection (95% granulocytes). Negative on peritoneal cells 4 days after thioglycollate injection. Negative on isolated lymphocytes, monocytes, erythrocytes and platelets.

Specificity:

Tissue staining:

RK-4 reaction pattern:

Tissues tested	Results
Bone marrow	positive on granulocytes positive on granulocyte precursors
	positive on segmented cells
	positive on band cells positive on metamyelocytes
	positive on myelocytes
Lung	negative on alveolar macrophages; interstitial tissues are occasionally positive.
Kidney	negative; on perivascular granulocytes occasionally positive.
Skin	negative; on perivascular granulocytes occasionally positive.
Liver	negative; on perivascular granulocytes occasionally positive.
Blood vessel wall	negative

Selected references

STEINHAUSEN: F;, H. Xu, H.-J. Groene, H.J. Peters: A monoclonal Antibody Selectively Recognizing Rat Granulocytes. Immunobiol: 181: 324 (1990)

For *in vitro* research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.