



T-4290

Rabbit anti Leu-Enkephalin

Leu-enkephalin is an endogenous opioid peptide neurotransmitter that is found naturally in the brains of many animals, including humans. It is one of the two forms of enkephalin; the other is met-enkephalin. The tyrosine residue at position 1 is thought to be analogous to the 3-hydroxyl group on morphine. Leu-enkephalin has agonistic actions at both the μ - and δ -opioid receptors, with significantly greater preference for the latter.

This antibody was generated by immunization of rabbits with Leu-Enkephalin coupled to a carrier protein.

TECHNICAL AND ANALYTICAL CHARACTERISTICS	
Lot number:	A07690
Host species:	Rabbit IgG
Quantity:	50µl
Format:	Neat undiluted antiserum, lyophilized, packaged under nitrogen. Reconstitute by adding 50µl distilled water. This will give the equivalent of undiluted antiserum.
Stability:	Original vial: at least one year at 4° - 8°C from date of delivery. Minimize repeated thawing and freezing of the antiserum by freezing aliquots at - 20°C or below.
Applications:	This antibody has been tested and validated in ELISA against Leu- Enkephalin. Other applications like immunohistochemistry (IHC), FACS or Western Blot may work as well. Optimal dilutions should be determined by the end user. Please see www.bma.ch for protocols and general information.
Immunogen:	Synthetic peptide H-Tyr-Gly-Gly-Phe-Leu-NH2 coupled to carrier protein.
Related Products:	S-1419: Met-Enkephalin ELISA, high sensitivity for extracted samples S-1541: Leu-Enkephalin ELISA, high sensitivity for extracted samples T-4288: rabbit anti Leu-Enkephalin, diluted antiserum T-4289: rabbit anti Leu-Enkephalin, purified IgG T-4292: rabbit anti Met-Enkephalin, diluted antiserum T-4297: rabbit anti Met-Enkephalin-Arg-Gly-Leu, purified IgG T-4298: rabbit anti Met-Enkephalin-Arg-Gly-Leu, neat antiserum T-4302: rabbit anti Met-Enkephalin-Arg-Phe, neat antiserum

This product contains no preservative and is intended for laboratory use and research purposes only. Purchase of this product does not include authorization to use it in diagnostic or therapeutic applications.