



BMA BIOMEDICALS



Peninsula Laboratories

---

## T-4298

### Rabbit anti Methionine Enkephalin-Arg-Gly-Leu

Met-Enkephalin is an endogenous opioid pentapeptide that has opioid effects of a relatively short duration. It is one of the two forms of enkephalin, the other being leu-enkephalin. The enkephalins are considered to be the primary endogenous ligands of the  $\delta$ -opioid receptor, due to their high potency and selectivity for the site over the other endogenous opioids. Met-enkephalin is found mainly in the adrenal medulla and throughout the central nervous system (CNS). It is synthesized from proenkephalin via proteolytic cleavage. As a result, its cleavage generates: the met-enkephalin derivatives, the heptapeptide met-enkephalin-arg-phe (261–267), and the octapeptide met-enkephalin-arg-gly-leu (186–193).

This antibody was generated by immunization of rabbits with Methionine Enkephalin-Arg-Gly-Leu coupled to a carrier protein.

---

#### TECHNICAL AND ANALYTICAL CHARACTERISTICS

<b>Lot number:</b>	022532
<b>Host species:</b>	Rabbit IgG
<b>Quantity:</b>	50 $\mu$ l
<b>Format:</b>	Neat undiluted antiserum, lyophilized, packaged under nitrogen. Reconstitute by adding 50 $\mu$ l distilled water. This will give the equivalent of undiluted antiserum.
<b>Stability:</b>	Original vial: at least one year at 4 $^{\circ}$ - 8 $^{\circ}$ C from date of delivery. Minimize repeated thawing and freezing of the antiserum by freezing aliquots at -20 $^{\circ}$ C or below.
<b>Applications:</b>	This antibody has been tested and validated in ELISA against Methionine Enkephalin-Arg-Gly-Leu. Other applications like immunohistochemistry (IHC), FACS or Western Blot may work as well. Optimal dilutions should be determined by the end user. Please see <a href="http://www.bma.ch">www.bma.ch</a> for protocols and general information.
<b>Immunogen:</b>	Synthetic peptide H-Tyr-Gly-Gly-Phe-Met-Arg-Gly-Leu-NH <sub>2</sub> coupled to carrier protein.

Related Products: S-1419: Met-Enkephalin ELISA, high sensitivity for extracted samples  
S-1541: Leu-Enkephalin ELISA, high sensitivity for extracted samples  
T-4288: rabbit anti Leu-Enkephalin, diluted antiserum  
T-4289: rabbit anti Leu-Enkephalin, purified IgG  
T-4290: rabbit anti Leu-Enkephalin, neat antiserum  
T-4292: rabbit anti Met-Enkephalin, diluted antiserum  
T-4297: rabbit anti Met-Enkephalin-Arg-Gly-Leu, purified IgG  
T-4302: rabbit anti Met-Enkephalin-Arg-Phe, neat antiserum  
T-4301: rabbit anti Met-Enkephalin-Arg-Phe, purified IgG

---

This product contains no preservative and is intended for laboratory use and research purposes only. Purchase of this product does not include authorization to use it in diagnostic or therapeutic applications.

T-4298

neat serum

1.3.2021